



TOBACCO USE

Healthy People 2020 Leading Health Indicators (LHIs)

Goal: Reduce illness, disability, and death related to tobacco use and secondhand smoke exposure.

TU-1.1 Reduce cigarette smoking by adults

BASELINE
20.6%

2008

17.0%

2014



of **adults** aged 18 years and older were current cigarette smokers in 2014

*(age adjusted to the year 2000 standard population)
Data Source: National Health Interview Survey*

TARGET
12%
YEAR 2020



TU-2.2 Reduce use of cigarettes by adolescents (past month)

BASELINE
19.5%

2008

15.7%

2013



of **adolescents** in grades 9 through 12 smoked cigarettes in the past 30 days in 2013

Data Source: Youth Risk Behavior Surveillance System

TARGET
16%
YEAR 2020



TU-11.1 Reduce the proportion of children aged 3 to 11 years exposed to secondhand smoke

BASELINE
52.2%

2005-2008

41.3%

2009-2012



of **children** aged 3 to 11 years were exposed to secondhand smoke in 2009 - 2012

Data Source: National Health and Nutrition Examination Survey

TARGET
47%
YEAR 2020



For more information, please visit <http://www.cdc.gov/tobacco/hp2020>

Email questions to: hp2020tobacco@cdc.gov



USE YOUR
SMARTPHONE
TO SCAN THIS
QR CODE



**Centers for Disease
Control and Prevention**
National Center for Chronic
Disease Prevention and
Health Promotion